arXiv:0812.0483v1 [gr-qc] 2 Dec 2008

Primordial production of massive relic gravitational waves from a weak modification of General Relativity

Christian Corda

December 2, 2008

Centro di Scienze Naturali, Via di Galceti 74 - 59100 PRATO, Italy and
Associazione Galileo Galilei, Via Pier Cironi 16 - 59100 PRATO, Italy

E-mail address: christian.corda@ego-gw.it

Abstract

The production of a stochastic background of relic gravitational waves is well known in various works in the literature, where, using the so called adiabatically-amplified zero-point fluctuations process it has been shown how the standard inflationary scenario for the early universe can in principle provide a distinctive spectrum of relic gravitational waves. In this paper, it is shown that a weak modification of General Relativity produces a third massive polarization of gravitational waves and the primordial production of this polarization is analysed adapting the adiabatically-amplified zero-point fluctuations process at this case.

The presence of the mass could also have important applications in cosmology as the fact that gravitational waves can have mass could give a contribution to the dark matter of the Universe.

At the end of the paper an upper bound for these relic gravitational waves, which arises from the WMAP constrains, is also released.

1 Introduction

Recently, the data analysis of interferometric gravitational waves (GWs) detectors has been started (for the current status of GWs interferometers see [1, 2, 3, 4, 5, 6, 7, 8]) and the scientific community aims in a first direct detection of GWs in next years.

Detectors for GWs will be important for a better knowledge of the Universe and also to confirm or ruling out the physical consistency of General Relativity or of any other theory of gravitation [9, 10, 11, 12, 13, 14]. This is because, in the context of Extended Theories of Gravity, some differences between General Relativity and the others theories can be pointed out starting by the linearized theory of gravity [9, 10, 12, 14]. In this picture, detectors for GWs are in principle sensitive also to a hypotetical *scalar* component of gravitational radiation, that appears in extended theories of gravity like scalar-tensor gravity and high order theories [12, 15, 16, 17, 18, 19, 20, 21, 22], Brans-Dicke theory [23] and string theory [24].

A possible target of these experiments is the so called stochastic background of gravitational waves [25, 26, 27, 28, 29, 30].

The production of the primordial part of this stochastick background (relic GWs) is well known in the literature starting by the works of [25, 26] and [27, 28], that, using the so called adiabatically-amplified zero-point fluctuations process, have shown in two different ways how the standard inflationary scenario for the early universe can in principle provide a distinctive spectrum of relic gravitational waves, while in [29, 30] the primordial production has been analyzed for the scalar component admitted from scalar-tensor gravity. In this paper, it is shown that a weak modification of General Relativity generates a third massive polarization of gravitational waves and the primordial production of this polarization is analysed adapting the adiabatically-amplified zero-point fluctuations process at this case. We have also to emphasize that, in a recent paper [40], such a process has been applied to the same theory which we are going to discuss in the present work. But, in [40] a different point of view has been considered. In that case, using a conform analysis, the authors discussed such a process in respect to the two standard polarizations which arises from standard General Relativity. In the present paper the analysis is focused to the third massive polarization.

The presence of the mass could also have important applications in cosmology because the fact that gravitational waves can have mass could give a contribution to the dark matter of the Universe.

At the end of the paper an upper bound for these relic gravitational waves, which arises from the WMAP constrains, is also released.

2 A weak modification of general relativity

Let us consider the action

$$S=\int d^4x\sqrt{-g}f_0R^{1+\varepsilon}+\mathcal{L}_m \tag{1}$$

Equation (1) is a particular choice in $f(R)$ theories of gravity [9, 10, 11, 13, 19, 21] in respect to the well known canonical one of General Relativity (the Einstein - Hilbert action [31, 32]) which is

$$S=\int d^4x\sqrt{-g}R+\mathcal{L}_m. \tag{2}$$

Criticisms on $f(R)$ theories of gravity arises from the fact that lots of such theories can be excluded by requirements of Cosmology and Solar System tests

[33]. But, in the case of the action (1), the variation from standard General Relativity is very weak, becauese ε is a very small real parameter, thus, the mentioned constrains could be, in principle, satisfed [33]. Note: General Relativity is obtained for $\varepsilon = 0$ and $f_0 = 1$.

The action (1) has been analyzed in [34] in a cosmologic context. But, because we will interact with gravitational waves, i.e. the linearized theory in vacuum, $\mathcal{L}_m = 0$ will be put and the pure curvature action

$$S = \int d^4x\sqrt{-g}f_0R^{1+\varepsilon} \tag{3}$$

will be considered.

3 The field equations

Following [32, 35] (note that in this paper we work with $8\pi G = 1$, $c = 1$ and $\hbar = 1$), the variational principle

$$\delta\int d^4x\sqrt{-g}f_0R^{1+\varepsilon} = 0 \tag{4}$$

in a local Lorentz frame will be used.

One gets:

$$\begin{array}{c} \delta\int d^4x\sqrt{-g}f_0R^{1+\varepsilon} = \int d^4x[\delta\sqrt{-g}f_0R^{1+\varepsilon} + f_0\sqrt{-g}\delta R^{1+\varepsilon}] = \\ \\ = \int d^4x[\sqrt{-g}f_0(1+\varepsilon)R^{\varepsilon}R_{\mu\nu} - \frac{1}{2}g_{\mu\nu}f_0R^{1+\varepsilon}]\delta g^{\mu\nu} + \\ \\ +\int d^4x\sqrt{-g}(1+\varepsilon)f_0R^{\varepsilon}g^{\mu\nu}\delta R_{\mu\nu}. \end{array} \tag{5}$$

Recalling the relation between the Christoffel coefficients and the Ricci tensor [32, 35] one can write

$$g^{\mu\nu}\delta R_{\mu\nu} = g^{\mu\nu}\partial_{\alpha}(\delta\Gamma^{\alpha}_{\mu\nu}) - g^{\mu\alpha}\partial_{\alpha}(\delta\Gamma^{\nu}_{\mu\nu}) \equiv \partial_{\alpha}X^{\alpha}, \tag{6}$$

where

$$X^{\alpha} \equiv g^{\mu\nu}(\delta\Gamma^{\alpha}_{\mu\nu}) - g^{\mu\alpha}(\delta\Gamma^{\nu}_{\mu\nu}). \tag{7}$$

In this way, the second integral in equation (5) can be computed as

$$\begin{array}{c} \int d^4x\sqrt{-g}(1+\varepsilon)f_0R^{\varepsilon}g^{\mu\nu}\delta R_{\mu\nu} = \int d^4x\sqrt{-g}(1+\varepsilon)f_0R^{\varepsilon}\partial_{\alpha}X^{\alpha} = \\ \\ = \int d^4x\partial_{\alpha}[\sqrt{-g}(1+\varepsilon)f_0R^{\varepsilon}X^{\alpha}] - \int d^4x\partial_{\alpha}[\sqrt{-g}(1+\varepsilon)f_0R^{\varepsilon}]X^{\alpha}. \end{array} \tag{8}$$

Assuming that fields are equal to zero at infinity [32, 35], one gets

$$d^4x\sqrt{-g}(1+\varepsilon)f_0R^{\varepsilon}g^{\mu\nu}\delta R_{\mu\nu} = -\int d^4x\partial_{\alpha}[\sqrt{-g}((1+\varepsilon)f_0R^{\varepsilon}]X^{\alpha}. \tag{9}$$

Now, let us compute X^{α}. Recalling that in a local Lorentz frame it is

$$\bigtriangledown_{\beta}\, g_{\mu\nu}=\partial_{\beta}g_{\mu\nu}=0 \tag{10}$$

and using the well known definitions of the Christofell coefficients [32, 35] it is

$$\begin{aligned}\delta\Gamma_{\mu\nu}^{\alpha}&=\delta[\tfrac{1}{2}g^{\beta\alpha}(\partial_{\mu}g_{\beta\nu}+\partial_{\nu}g_{\mu\beta}-\partial_{\beta}g_{\mu\nu})]=\\&=\tfrac{1}{2}g^{\beta\alpha}(\partial_{\mu}\delta g_{\beta\nu}+\partial_{\nu}\delta g_{\mu\beta}-\partial_{\beta}\delta g_{\mu\nu}).\end{aligned} \tag{11}$$

In the same way it is

$$\delta\Gamma_{\mu\nu}^{\nu}=\frac{1}{2}g^{\nu\beta}\partial_{\mu}(\delta g_{\nu\beta}). \tag{12}$$

From eqs. (11) and (12) one gets

$$g^{\mu\nu}(\delta\Gamma_{\mu\nu}^{\alpha})=\frac{1}{2}\partial^{\alpha}(g_{\mu\nu}\delta g^{\mu\nu})-\partial^{\mu}(g_{\beta\mu}\delta g^{\nu\beta}) \tag{13}$$

and

$$g^{\mu\alpha}(\delta\Gamma_{\mu\nu}^{\nu})=-\frac{1}{2}\partial^{\alpha}(g_{\nu\beta}\delta g^{\nu\beta}). \tag{14}$$

Then, substituting in (7), it is

$$X^{\alpha}=\partial^{\alpha}(g_{\mu\nu}\delta g^{\mu\nu})-\partial^{\mu}(g_{\mu\nu}\delta g^{\alpha\nu}). \tag{15}$$

With this equation, equation (9) becomes

$$\begin{aligned}&\textstyle\int d^{4}x\sqrt{-g}(1+\varepsilon)f_{0}R^{\varepsilon}g^{\mu\nu}\delta R_{\mu\nu}=\\&=\textstyle\int d^{4}x\partial_{\alpha}[\sqrt{-g}(1+\varepsilon)f_{0}R^{\varepsilon}][\partial^{\mu}(g_{\mu\nu}\delta g^{\alpha\nu})-\partial^{\alpha}(g_{\mu\nu}\delta g^{\mu\nu})],\end{aligned} \tag{16}$$

which also gives

$$\begin{aligned}&\textstyle\int d^{4}x\sqrt{-g}(1+\varepsilon)f_{0}R^{\varepsilon}g^{\mu\nu}\delta R_{\mu\nu}=\\&=\textstyle\int d^{4}x\{g_{\mu\nu}\partial^{\alpha}\partial_{\alpha}[\sqrt{-g}(1+\varepsilon)f_{0}R^{\varepsilon}]\delta g^{\mu\nu}\}-\int d^{4}x\{g_{\mu\nu}\partial^{\mu}\partial_{\alpha}[\sqrt{-g}(1+\varepsilon)f_{0}R^{\varepsilon}]\delta g^{\alpha\nu}\}.\end{aligned} \tag{17}$$

Putting this equation in the variation (5) one obtains

$$\begin{aligned}&\delta\textstyle\int d^{4}x\sqrt{-g}(1+\varepsilon)f_{0}R^{\varepsilon}=\int d^{4}x[\sqrt{-g}(1+\varepsilon)f_{0}R^{\varepsilon}R_{\mu\nu}-\frac{1}{2}g_{\mu\nu}f_{0}R^{1+\varepsilon}]\delta g^{\mu\nu}+\\&+\textstyle\int d^{4}x\{g_{\mu\nu}\partial^{\alpha}\partial_{\alpha}[\sqrt{-g}(1+\varepsilon)f_{0}R^{\varepsilon})]-g_{\alpha\nu}\partial^{\mu}\partial_{\alpha}[\sqrt{-g}(1+\varepsilon)f_{0}R^{\varepsilon}]\}\delta g^{\mu\nu}\}.\end{aligned} \tag{18}$$

The above variation is equal to zero for

$$(1+\varepsilon)f_{0}R^{\varepsilon}R_{\mu\nu}-\frac{1}{2}g_{\mu\nu}f_{0}R^{1+\varepsilon}=(\bigtriangledown_{\mu}\bigtriangledown_{\nu}-g_{\mu\nu}\Box)(1+\varepsilon)f_{0}R^{\varepsilon}, \tag{19}$$

which are the modified Einstein field equations. Writing down, exlplicitly, the Einstein tensor eqs. (19) become

$$G_{\mu\nu}=\frac{1}{(1+\varepsilon)f_0R^{\varepsilon}}\{-\frac{1}{2}g_{\mu\nu}\varepsilon f_0R^{1+\varepsilon}+[(1+\varepsilon)f_0R^{\varepsilon}]_{;\mu;\nu}-g_{\mu\nu}\Box[(1+\varepsilon)f_0R^{\varepsilon}]\}. \tag{20}$$

Taking the trace of the field equations (20) one gets

$$\Box(1+\varepsilon)f_0R^{\varepsilon}=\frac{(1-\varepsilon)}{3}f_0R^{1+\varepsilon}. \tag{21}$$

Now, we can define the *effective* scalar field

$$\Phi\equiv(1+\varepsilon)f_0R^{\varepsilon} \tag{22}$$

with associated an effective potential

$$\frac{dV}{d\Phi}\equiv\frac{(1-\varepsilon)}{3}f_0R^{1+\varepsilon}. \tag{23}$$

Thus, from eq. (21), a Klein - Gordon equation for the effective Φ scalar field is obtained:

$$\Box\Phi=\frac{dV}{d\Phi}. \tag{24}$$

4 The linearized theory

To study gravitational waves, the linearized theory has to be analyzed, with a little perturbation of the background, which is assumed given by a near Minkowskian background, i.e. a Minkowskian background plus $\Phi=\Phi_0$ (the Ricci scalar is assumed constant in the background) [9, 19]. We also assume Φ_0 to be a minimum for the effective potential V:

$$V\simeq\frac{1}{2}\alpha\delta\Phi^2\Rightarrow\frac{dV}{d\Phi}\simeq m^2\delta\Phi, \tag{25}$$

and the constant m has mass dimension.

Putting

$$\begin{array}{c} g_{\mu\nu}=\eta_{\mu\nu}+h_{\mu\nu}\\ \\ \Phi=\Phi_0+\delta\Phi. \end{array} \tag{26}$$

to first order in $h_{\mu\nu}$ and $\delta\Phi$, calling $\widetilde{R}_{\mu\nu\rho\sigma}$, $\widetilde{R}_{\mu\nu}$ and $\widetilde{R}$ the linearized quantity which correspond to $R_{\mu\nu\rho\sigma}$, $R_{\mu\nu}$ and R, the linearized field equations are obtained [12, 19, 31]:

$$\begin{array}{c} \widetilde{R}_{\mu\nu}-\frac{\widetilde{R}}{2}\eta_{\mu\nu}=(\partial_\mu\partial_\nu h_m-\eta_{\mu\nu}\Box h_m)\\ \\ \Box h_m=m^2h_m, \end{array} \tag{27}$$

where

$$h_m \equiv \frac{\delta\Phi}{\Phi_0}. \tag{28}$$

Then, from the second of eqs. (27), one can define the mass like

$$m \equiv \sqrt{\frac{\Box h_m}{h_m}} = \sqrt{\frac{\Box \delta\Phi}{\delta\Phi}} = \sqrt{\frac{\Box \delta R^{\varepsilon}}{\delta R^{\varepsilon}}}. \tag{29}$$

Thus, as the mass is generated by variation of the Ricci scalar, we can say that, in a certain sense, it is generated by variation of spacetime curvature, re-obtaining the same result of [9, 19]. The difference with the works [9, 19] is that now the theory is more suitable as the modification of General Relativity is very weak and in agreement with requirements of Cosmology and Solar System tests [33].

$\widetilde{R}_{\mu\nu\rho\sigma}$ and eqs. (27) are invariants for gauge transformations [9, 12, 19]

$$\begin{array}{c} h_{\mu\nu} \rightarrow h'_{\mu\nu} = h_{\mu\nu} - \partial_{(\mu}\epsilon_{\nu)} \\ \\ \delta\Phi \rightarrow \delta\Phi' = \delta\Phi; \end{array} \tag{30}$$

then

$$\bar{h}_{\mu\nu} \equiv h_{\mu\nu} - \frac{h}{2}\eta_{\mu\nu} + \eta_{\mu\nu} h_m \tag{31}$$

can be defined, and, considering the transform for the parameter ϵ^{μ}

$$\Box \epsilon_{\nu} = \partial^{\mu} \bar{h}_{\mu\nu}, \tag{32}$$

a gauge parallel to the Lorenz one of electromagnetic waves can be choosen:

$$\partial^{\mu} \bar{h}_{\mu\nu} = 0. \tag{33}$$

In this way field equations read like

$$\Box \bar{h}_{\mu\nu} = 0 \tag{34}$$

$$\Box h_m = m^2 h_m \tag{35}$$

Solutions of eqs. (34) and (35) are plan waves [12, 19]:

$$\bar{h}_{\mu\nu} = A_{\mu\nu}(\overrightarrow{p}) \exp(ip^{\alpha}x_{\alpha}) + c.c. \tag{36}$$

$$h_m = a(\overrightarrow{p}) \exp(iq^{\alpha}x_{\alpha}) + c.c. \tag{37}$$

where

$$
\begin{array}{ll}
k^{\alpha} \equiv (\omega, \overrightarrow{p}) & \omega = p \equiv |\overrightarrow{p}| \\
\\
q^{\alpha} \equiv (\omega_m, \overrightarrow{p}) & \omega_m = \sqrt{m^2 + p^2}.
\end{array}
\tag{38}
$$

In eqs. (34) and (36) the equation and the solution for the standard waves of General Relativity [31, 32] have been obtained, while eqs. (35) and (37) are respectively the equation and the solution for the massive mode (see also [9, 12, 19]).

The fact that the dispersion law for the modes of the massive field h_m is not linear has to be emphatized. The velocity of every "ordinary" (i.e. which arises from General Relativity) mode $\bar{h}_{\mu\nu}$ is the light speed c, but the dispersion law (the second of eq. (38)) for the modes of h_m is that of a massive field which can be discussed like a wave-packet [9, 12, 19]. Also, the group-velocity of a wave-packet of h_m centered in $\overrightarrow{p}$ is

$$
\overrightarrow{v_G} = \frac{\overrightarrow{p}}{\omega}, \tag{39}
$$

which is exactly the velocity of a massive particle with mass m and momentum $\overrightarrow{p}$.

From the second of eqs. (38) and eq. (39) it is simple to obtain:

$$
v_G = \frac{\sqrt{\omega^2 - m^2}}{\omega}. \tag{40}
$$

Then, wanting a constant speed of the wave-packet, it has to be [9, 12, 19]

$$
m = \sqrt{(1 - v_G^2)}\omega. \tag{41}
$$

The relation (41) is shown in fig. 1 for a value $v_G = 0.9$.

Now, the analisys can remain in the Lorenz gauge with trasformations of the type $\Box\epsilon_\nu = 0$; this gauge gives a condition of transversality for the ordinary part of the field: $k^\mu A_{\mu\nu} = 0$, but does not give the transversality for the total field $h_{\mu\nu}$. From eq. (31) it is

$$
h_{\mu\nu} = \bar{h}_{\mu\nu} - \frac{\bar{h}}{2}\eta_{\mu\nu} + \eta_{\mu\nu} h_m. \tag{42}
$$

At this point, if being in the massless case [9, 12, 19], it could been put

$$
\begin{array}{c}
\Box\epsilon^\mu = 0 \\
\\
\partial_\mu \epsilon^\mu = -\frac{\bar{h}}{2} + h_m,
\end{array}
\tag{43}
$$

which gives the total transversality of the field. But in the massive case this is impossible. In fact, applying the Dalembertian operator to the second of eqs. (43) and using the field equations (34) and (35) it results

$$
\Box\epsilon^\mu = m^2 h_m, \tag{44}
$$



Figure 1: the mass-frequency relation for a massive GW propagating with a speed of $0.9c$: for the mass it is $1Hz = 10^{-15}eV$

which is in contrast with the first of eqs. (43). In the same way it is possible to show that it does not exist any linear relation between the tensorial field $\bar{h}_{\mu\nu}$ and the massive field h_m. Thus a gauge in wich $h_{\mu\nu}$ is purely spatial cannot be chosen (i.e. it cannot be put $h_{\mu 0}=0$, see eq. (42)) . But the traceless condition to the field $\bar{h}_{\mu\nu}$ can be put :

$$\begin{array}{c} \square\epsilon^{\mu}=0 \\ \\ \partial_{\mu}\epsilon^{\mu}=-\frac{\bar{h}}{2}. \end{array} \tag{45}$$

These equations imply

$$\partial^{\mu}\bar{h}_{\mu\nu}=0. \tag{46}$$

To save the conditions $\partial_{\mu}\bar{h}^{\mu\nu}$ and $\bar{h}=0$ transformations like

$$\begin{array}{c} \square\epsilon^{\mu}=0 \\ \\ \partial_{\mu}\epsilon^{\mu}=0 \end{array} \tag{47}$$

can be used and, taking $\overrightarrow{p}$ in the z direction, a gauge in which only A_{11}, A_{22}, and $A_{12}=A_{21}$ are different to zero can be chosen. The condition $\bar{h}=0$ gives $A_{11}=-A_{22}$. Now, putting these equations in eq. (42), it results

$$h_{\mu\nu}(t,z)=A^{+}(t-z)e_{\mu\nu}^{(+)}+A^{\times}(t-z)e_{\mu\nu}^{(\times)}+h_m(t-v_G z)\eta_{\mu\nu}. \tag{48}$$

The term $A^{+}(t-z)e_{\mu\nu}^{(+)}+A^{\times}(t-z)e_{\mu\nu}^{(\times)}$ describes the two standard polarizations of gravitational waves which arise from General Relativity, while the term $h_m(t-v_G z)\eta_{\mu\nu}$ is the massive field arising from the high order theory. In other words, the function R^{ε} of the Ricci scalar generates a third massive polarization for gravitational waves which is not present in standard General Relativity.

5 The primordial production of the third polarization

Now, let us consider the primordial physical process, which gave rise to a characteristic spectrum Ω_{gw} for the relic GWs. Such physical process has been analyzed in different ways: respectively in refs. [25, 26] and [27, 28] but only for the components of eq. (48) which arises from General Relativity, while in [29] the process has been extended to scalar-tensor gravity. Actually the process can be furtherly improved showing the primordial production of the third polarization of eq. (48).

Before starting with the analysis, it has to be emphasized that, considering a stochastic background of GWs, it can be characterized by a dimensionless spectrum [25, 26, 27, 28, 29]

$$\Omega_{gw}(f)\equiv\frac{1}{\rho_c}\frac{d\rho_{gw}}{d\ln f}, \tag{49}$$

where

$$\rho_c \equiv \frac{3H_0^2}{8G} \tag{50}$$

is the (actual) critical density energy, ρ_c of the Universe, H_0 the actual value of the Hubble expansion rate and $d\rho_{gw}$ the energy density of relic GWs in the frequency range f to $f + df$.

The existence of a relic stochastic background of GWs is a consequence of generals assumptions. Essentially it derives from a mixing between basic principles of classical theories of gravity and of quantum field theory. The strong variations of the gravitational field in the early universe amplifie the zero-point quantum oscillations and produce relic GWs. It is well known that the detection of relic GWs is the only way to learn about the evolution of the very early universe, up to the bounds of the Planck epoch and the initial singularity [21, 25, 26, 27, 28, 29]. It is very important to stress the unavoidable and fundamental character of this mechanism. The model derives from the inflationary scenario for the early universe [36, 37], which is tuned in a good way with the WMAP data on the Cosmic Background Radiation (CBR) (in particular exponential inflation and spectral index ≈ 1 [38, 39]). Inflationary models of the early Universe were analysed in the early and middles 1980's (see [36] for a review), starting from an idea of A. Guth [37]. These are cosmological models in which the Universe undergoes a brief phase of a very rapid expansion in early times. In this context the expansion could be power-law or exponential in time. Inflationary models provide solutions to the horizon and flatness problems and contain a mechanism which creates perturbations in all fields. Important for our goals is that this mechanism also provides a distinctive spectrum of relic GWs. The GWs perturbations arise from the uncertainty principle and the spectrum of relic GWs is generated from the adiabatically-amplified zero-point fluctuations [21, 25, 26, 27, 28, 29].

Now, the calculation for a simple inflationary model will be shown for the third polarization of eq. (48), following the works of Allen [25, 26] that performed the calculation in the case of standard General Relativity and Corda, Capozziello and De Laurentis [29, 30] that extended the process to scalar GWs. Even here we have to recall that, in a recent paper [40], such a process has been applied to the theory arising from the action (1). But, in [40] a different point of view has been considered. In that case, using a conform analysis, the authors discussed such a process in respect to the two standard polarizations which arises from standard General Relativity. In the following the analysis is focused to the third massive polarization. Thus, in a certain sense, one can say that the present analysis is an integration of the analysis in [40].

It will be assumed that the universe is described by a simple cosmology in two stages, an inflationary De Sitter phase and a radiation dominated phase [21, 25, 26, 27, 28, 29]. The line element of the spacetime is given by

$$ds^2 = a^2(\eta)[-d\eta^2 + d\overrightarrow{x}^2 + h_{\mu\nu}(\eta, \overrightarrow{x})dx^\mu dx^\nu]. \tag{51}$$

In this line element, because we are considering only the third polarization, the metric perturbation (48) reduces to

$$h_{\mu\nu} = h_m I_{\mu\nu}, \tag{52}$$

where

$$I_{\mu\nu} \equiv \begin{matrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1. \end{matrix} \tag{53}$$

In the De Sitter phase ($\eta < \eta_1$) the equation of state is $P = -\rho = const$, the scale factor is $a(\eta) = \eta_1^2 \eta_0^{-1}(2\eta_1 - \eta)^{-1}$ and the Hubble constant is given by $H(\eta) = H_{ds} = c\eta_0/\eta_1^2$.

In the radiation dominated phase ($\eta > \eta_1$) the equation of state is $P = \rho/3$, the scale factor is $a(\eta) = \eta/\eta_0$ and the Hubble constant is given by $H(\eta) = c\eta_0/\eta^2$ [21, 25, 26, 29, 30].

Expressing the scale factor in terms of comoving time defined by

$$cdt = a(t)d\eta \tag{54}$$

one gets

$$a(t) \propto \exp(H_{ds}t) \tag{55}$$

during the De Sitter phase and

$$a(t) \propto \sqrt{t} \tag{56}$$

during the radiation dominated phase. In order to obtain a solution for the horizon and flatness problems it has to be [36, 37]

$$\tfrac{a(\eta_0)}{a(\eta_1)} > 10^{27}$$

The third polarization generates weak perturbations $h_{\mu\nu}(\eta, \overrightarrow{x})$ of the metric (52) that can be written in the form

$$h_{\mu\nu} = I_{\mu\nu}(\hat{k})X(\eta)\exp(\overrightarrow{k} \cdot \overrightarrow{x}), \tag{57}$$

in terms of the conformal time η where $\overrightarrow{k}$ is a constant wavevector and

$$h_m(\eta, \overrightarrow{k}, \overrightarrow{x}) = X(\eta)\exp(\overrightarrow{k} \cdot \overrightarrow{x}). \tag{58}$$

By putting $Y(\eta) = a(\eta)X(\eta)$ and with the standard linearized calculation in which the connections (i.e. the Cristoffel coefficents), the Riemann tensor, the Ricci tensor and the Ricci scalar curvature are found, from Friedman linearized equations it is obtained that the function $Y(\eta)$ satisfies the equation

$$Y'' + (|\overrightarrow{k}|^2 - \frac{a''}{a})Y = 0 \tag{59}$$

where $'$ denotes derivative with respect to the conformal time. Cleary, this is the equation for a parametrically disturbed oscillator.

The solutions of eq. (59) give us the solutions for the function $X(\eta)$, that can be expressed in terms of elementary functions simple cases of half integer Bessel or Hankel functions [21, 25, 26, 29, 30] in both the inflationary and radiation dominated eras:

For $\eta < \eta_1$

$$X(\eta) = \frac{a(\eta_1)}{a(\eta)}[1 + H_{ds}\omega^{-1}] \exp -ik(\eta - \eta_1), \tag{60}$$

for $\eta > \eta_1$

$$X(\eta) = \frac{a(\eta_1)}{a(\eta)}[\alpha \exp -ik(\eta - \eta_1) + \beta \exp ik(\eta - \eta_1), \tag{61}$$

where $\omega = ck/a$ is the angular frequency of the wave (that is function of the time because of the constance of $k = |\overrightarrow{k}|$), α and β are time-indipendent constants which can be obtained demanding that both X and $dX/d\eta$ are continuous at the boundary $\eta = \eta_1$ between the inflationary and the radiation dominated eras of the cosmologic expansion. With this constrain it is

$$\alpha = 1 + i\frac{\sqrt{H_{ds}H_0}}{\omega} - \frac{H_{ds}H_0}{2\omega^2} \tag{62}$$

$$\beta = \frac{H_{ds}H_0}{2\omega^2} \tag{63}$$

In eqs. (62), (63) $\omega = ck/a(\eta_0)$ is the angular frequency that would be observed today. Calculations like this are referred in the literature as Bogoliubov coefficient methods [21, 25, 26, 29, 30].

As inflation dampes out any classical or macroscopic perturbations, the minimum allowed level of fluctations is that requiered by the uncertainty principle. The solution (60) corresponds precisely to this De Sitter vacuum state [21, 25, 26, 29, 30]. Then, if the period of inflation was long enough, the observable properties of the Universe today should be the same properties of a Universe started in the De Sitter vacuum state.

In the radiation dominated phase the coefficients of α are the eigenmodes which describe particles while the coefficients of β are the eigenmodes which describe antiparticles. Thus, the number of created particles of angular frequency ω in this phase is

$$N_\omega = |\beta_\omega|^2 = (\frac{H_{ds}H_0}{2\omega^2})^2. \tag{64}$$

Now, one can write an expression for the energy spectrum of the relic gravitational waves background in the frequency interval $(\omega, \omega + d\omega)$ as

$$d\rho_{gw} = 2\hbar\omega(\frac{\omega^2 d\omega}{2\pi^2 c^3})N_\omega = \frac{\hbar H_{ds}^2 H_0^2}{4\pi^2 c^3}\frac{d\omega}{\omega} = \frac{\hbar H_{ds}^2 H_0^2}{4\pi^2 c^3}\frac{df}{f}. \tag{65}$$

Eq. (65) can be rewritten in terms of the present day and the De Sitter energy-density of the Universe. The Hubble expansion rates is

$$H_0^2 = \frac{8\pi G\rho_c}{3c^2},\ H_{ds}^2 = \frac{8\pi G\rho_{ds}}{3c^2}.$$

Then, defining the Planck density

$$\rho_{Planck} \equiv \frac{c^7}{\hbar G^2} \tag{66}$$

the spectrum is

$$\Omega_{gw}(f) = \frac{1}{\rho_c}\frac{d\rho_{sgw}}{d\ln f} = \frac{f}{\rho_c}\frac{d\rho_{gw}}{df} = \frac{16}{9}\frac{\rho_{ds}}{\rho_{Planck}}. \tag{67}$$

Some comments are needed. It has to be emphasized that the computation works for a very simplified model that does not include the matter dominated era. Including this era, the redshift has to be considered. An enlighting computation parallel to the one in [26] gives

$$\Omega_{gw}(f) = \frac{16}{9}\frac{\rho_{ds}}{\rho_{Planck}}(1+z_{eq})^{-1}, \tag{68}$$

for the waves which at the time in which the Universe was becoming matter dominated had a frequency higher than H_{eq}, the Hubble constant at that time. This corresponds to frequencies $f > (1+z_{eq})^{1/2}H_0$, where z_{eq} is the redshift of the Universe when the matter and radiation energy density were equal. The redshift correction in equation (68) is needed because the Hubble parameter, which is governed by Friedman equations, should be different from the observed one H_0 for a Universe without matter dominated era.

At lower frequencies the spectrum is [21, 25, 26, 29, 30]

$$\Omega_{gw}(f) \propto f^{-2}. \tag{69}$$

Moreover, let us note that the results (67) and (68), which are not frequency dependent, cannot be applied to all the frequencies. For waves with frequencies less than H_0 today, the energy density cannot be defined, because the wavelenght becomes longer than the Hubble radius. In the same way, at high frequencies there is a maximum frequency above which the spectrum drops to zero rapidly. In the above computation it has been implicitly assumed that the phase transition from the inflationary to the radiation dominated epoch is istantaneous. In the real Universe this phase transition occurs over some finite time $\Delta\tau$, and above a frequency

$$f_{max} = \frac{a(t_1)}{a(t_0)}\frac{1}{\Delta\tau}, \tag{70}$$

which is the redshifted rate of the transition, Ω_{gw} drops rapidly. These two cutoffs, at low and high frequencies, to the spectrum force the total energy density of the relic gravitational waves to be finite. For GUT energy-scale inflation it is [21, 25, 26, 29, 30].

$$\frac{\rho_{ds}}{\rho_{Planck}} \approx 10^{-12}. \tag{71}$$

6 Tuning with WMAP data

It is well known that WMAP observations put strongly severe restrictions on the spectrum of relic gravitational waves. In fig. 2 the spectrum $\Omega_{gw}$is mapped following [20]: the amplitude is chosen (determined by the ratio $\frac{\rho_{ds}}{\rho_{Planck}}$) to be *as large as possible, consistent with the WMAP constraints* on *tensor perturbations.* Nevertheless, because the spectrum falls off $\propto f^{-2}$ at low frequencies, this means that today, at LIGO-Virgo and LISA frequencies (indicate by the lines in fig. 2) [20], it is

$$\Omega_{gw}(f)h_{100}^2 < 9 * 10^{-13}. \tag{72}$$

It is interesting to calculate the correspondent strain at $\approx 100Hz$, where interferometers like Virgo and LIGO have a maximum in sensitivity. The well known equation for the characteristic amplitude, adapted for the third component of GWs can be used [20]:

$$h_{mc}(f) \simeq 1.26 * 10^{-18}(\frac{1Hz}{f})\sqrt{h_{100}^2\Omega_{gw}(f)}, \tag{73}$$

obtaning [20]

$$h_{mc}(100Hz) < 1.7 * 10^{-26}. \tag{74}$$

Then, as we expect a sensitivity of the order of 10^{-22} for our interferometers at $\approx 100Hz$, we need to gain four order of magnitude. Let us analyze smaller frequencies too. The sensitivity of the Virgo interferometer is of the order of 10^{-21} at $\approx 10Hz$ and in that case it is [20]

$$h_{mc}(10Hz) < 1.7 * 10^{-25}. \tag{75}$$

The sensitivity of the LISA interferometer will be of the order of 10^{-22} at $10^{-3} \approx Hz$ and in that case it is [20]

$$h_{mc}(100Hz) < 1.7 * 10^{-21}. \tag{76}$$

Then, a stochastic background of relic gravitational waves could be in principle detected by the LISA interferometer.

We emphasize that the assumption that all the tensorial perturbation in the Universe are due to a stochastic background of GWs is quit strong, but our results (72), (74), (75) and (76) can be considered like upper bounds.



Figure 2: adapted from C. Corda - Mod. Phys. Lett. A No. 22, 16, 1167-1173 (2007).
The spectrum of relic sGWs in inflationary models is flat over a wide range of frequencies. The horizontal axis is $\log_{10}$ of frequency, in Hz. The vertical axis is $\log_{10}\Omega_{gsw}$. The inflationary spectrum rises quickly at low frequencies (wave which rentered in the Hubble sphere after the Universe became matter dominated) and falls off above the (appropriately redshifted) frequency scale f_{max} associated with the fastest characteristic time of the phase transition at the end of inflation. The amplitude of the flat region depends only on the energy density during the inflationary stage; we have chosen the largest amplitude consistent with the WMAP constrains on scalar perturbations. This means that at LIGO and LISA frequencies, $\Omega_{gw}(f)h_{100}^2 < 9 * 10^{-13}$

7 Conclusions

It has been shown that a weak modification of general relativity produces a third massive polarization of gravitational waves and the primordial production of this polarization has been analysed adapting the adiabatically-amplified zero-point fluctuations process at this case.

The presence of the mass could also have important applications in cosmology because the fact that gravitational waves can have mass could give a contribution to the dark matter of the Universe.

At the end of the paper, an upper bound for these relic gravitational waves, which arises from the WMAP constrains, has also been released .

Acknowledgements

I would like to thank Salvatore Capozziello and Maria Felicia De Laurentis for helpful advices during my work. The European Gravitational Observatory (EGO) consortium has also to be thanked for the using of computing facilities.

References

[1] F. Acernese et al. (the Virgo Collaboration) - Class. Quant. Grav. **24,** 19, S381-S388 (2007)

[2] C. Corda - Astropart. Phys. **27,** No 6, 539-549 (2007);

[3] C. Corda - Int. J. Mod. Phys. D **16,** 9, 1497-1517 (2007)

[4] B. Willke et al. - Class. Quant. Grav. **23** 8S207-S214 (2006)

[5] D. Sigg (for the LIGO Scientific Collaboration) - www.ligo.org/pdf_public/P050036.pdf

[6] B. Abbott et al. (the LIGO Scientific Collaboration) - Phys. Rev. D 72, 042002 (2005)

[7] M. Ando and the TAMA Collaboration - Class. Quant. Grav. **19** 7 1615-1621 (2002)

[8] D. Tatsumi, Tsunesada Y and the TAMA Collaboration - Class. Quant. Grav. **21** 5 S451-S456 (2004)

[9] C. Corda - J. Cosmol. Astropart. Phys. JCAP04009 (2007)

[10] C. Corda - Int. Journ. Mod. Phys. A 23, 10, 1521-1535 (2008)

[11] G. Allemandi, M. Francaviglia, M. L. Ruggiero and A. Tartaglia - Gen. Rel. Grav. 37 11 (2005)

[12] Capozziello S and C. Corda - Int. J. Mod. Phys. D **15,** 1119 -1150 (2006)

[13] C. Corda - *Response of laser interferometers to scalar gravitational waves*- talk in the *Gravitational Waves Data Analysis Workshop in the General Relativity Trimester of the Institut Henri Poincare* - Paris 13-17 November 2006, on the web in www.luth2.obspm.fr/IHP06/workshops/gwdata/corda.pdf

[14] C. Corda- Astropart. Phys. 28, 247-250 (2007)

[15] M. Shibata, K. Nakao and T. Nakamura - Phys. Rev. D **50**, 7304 (1994)

[16] M. Maggiore and A. Nicolis - Phys. Rev. D **62** 024004 (2000); also in gr-qc/9907055

[17] T. Damour and G. Esposito-Farese - Class. Quant. Grav. **9** 2093-2176 (1992); M. E. Tobar , T. Suzuki and K. Kuroda Phys. Rev. D 59 102002 (1999)

[18] K. Nakao, T. Harada , M. Shibata, S. Kawamura and T. Nakamura - Phys. Rev. D 63, 082001 (2001)

[19] C. Corda and M. F. De Laurentis - Proceedings of the 10th ICATPP Conference on Astroparticle, Particle, Space Physics, Detectors and Medical Physics - Applications, Villa Olmo, Como, Italy (October 8-12 2007)

[20] C. Corda - Mod. Phys. Lett. A No. 22, 16, 1167-1173 (2007)

[21] S. Capozziello, C. Corda and M. F. De Laurentis - Mod. Phys. Lett. A 22, 15, 1097-1104 (2007)

[22] C. Corda - Mod. Phys. Lett. A No. 22, 23, 1727-1735 (2007)

[23] C. Brans and R. H. Dicke - Phys. Rev. 124, 925 (1961)

[24] N. Bonasia and M. Gasperini - Phys. Rew. D **71** 104020 (2005)

[25] B. Allen - Proceedings of the Les Houches School on Astrophysical Sources of Gravitational Waves, eds. Jean-Alain Marck and Jean-Pierre Lasota (Cambridge University Press, Cambridge, England 1998).

[26] B. Allen - Phys. Rev. D **37**, 2078 (1988)

[27] L. P. Grishchuk and others - Phys. Usp. 44 1-51 (2001)

[28] L. P. Grishchuk and others - Usp. Fiz. Nauk 171 3 (2001)

[29] C. Corda, S. Capozziello and M. F. De Laurentis - AIP Conference Proceedings, Volume 966, pp 257-263 (2007) - Proceedings of the Fourth Italian-Sino Workshop on Relativistic Astrophysics July 20-30 2007, Pescara, Italy

[30] S. Capozziello, C. Corda and M. F. De Laurentis - Mod. Phys. Lett. A 22, 35, 2647-2655 (2007)

[31] C. W. Misner, K. S. Thorne and J. A. Wheeler - "Gravitation" - W.H.Feeman and Company - 1973

[32] L. Landau and E. Lifsits - "Teoria dei campi" - Editori riuniti edition III (1999)

[33] T. Clifton and J. D. Barrow - Phys. Rev. D **72,** 103005 (2005)

[34] S. Capozziello, E. Piedipalumbo and C. Rubano - Class. Quant. Grav. 23 1205 - 1216 (2006)

[35] S. Capozziello, and M. Funaro - “Introduzione alla Relatività Generale” - Liguori Editore Napoli, Italy (2005)

[36] G. Watson- “An exposition on inflationary cosmology” - North Carolina University Press (2000)

[37] A. Guth - Phys. Rev. **23** 347 (1981)

[38] C. L. Bennett et al - ApJS **148** 1 (2003)

[39] D.N. Spergel et al. - ApJS **148** 195 (2003)

[40] S. Capozziello, M. F. De Laurentis and M. Francaviglia - Astropart. Phys. **2,** No 2, 125-129 (2008)